arXiv:0704.2331v1 [math.AG] 18 Apr 2007

SYMMETRIES IN THE SYSTEM OF TYPE $D_4^{(1)}$

YUSUKE SASANO

Abstract. In this paper, we propose a 4-parameter family of coupled Painlevé III systems in dimension four with affine Weyl group symmetry of type $D_4^{(1)}$. We also propose its symmetric form in which the $D_4^{(1)}$-symmetries become clearly visible.

0. Statement of main results

In [9, 10, 11], we presented some types of coupled Painlevé systems with various affine Weyl group symmetries. In this paper, we present a 4-parameter family of coupled Painlevé III systems with affine Weyl group symmetry of type $D_4^{(1)}$. Since these universal Bäcklund transformations have Lie theoretic origin, similarity reduction of a Drinfeld-Sokolov hierarchy admits such a Bäcklund symmetry. The aim of this paper is to introduce the system of type $D_4^{(1)}$. After our discovery of this system, they were studied from the viewpoint of Drinfeld-Sokolov hierarchy by K. Fuji independently (cf. [1]), and he succeeded to obtain our system by similarity reduction of the Drinfeld-Sokolov hierarchy of type $D_4^{(1)}$.

At first, we propose a 4-parameter family of autonomous ordinary differential systems with the invariant divisors f_i as variables:

$$(1)\qquad \begin{cases} \dfrac{df_0}{dt}=-(2f_1g_1+\alpha_1)f_0-\alpha_0f_1,\\ \dfrac{df_1}{dt}=-(2f_0g_1+\alpha_0)f_1-\alpha_1f_0,\\ \dfrac{df_2}{dt}=\{(f_0+f_1)g_1+(f_3+f_4)g_2+1\}f_2-2\alpha_2g_1g_2,\\ \dfrac{df_3}{dt}=-(2f_4g_2+\alpha_4)f_3-\alpha_3f_4,\\ \dfrac{df_4}{dt}=-(2f_3g_2+\alpha_3)f_4-\alpha_4f_3,\\ \dfrac{dg_1}{dt}=(f_0+f_1)g_1^2-\{(f_3+f_4)g_2-\alpha_0-\alpha_1\}g_1+(f_3+f_4)f_2,\\ \dfrac{dg_2}{dt}=(f_3+f_4)g_2^2-\{(f_0+f_1)g_1-\alpha_3-\alpha_4\}g_2+(f_0+f_1)f_2. \end{cases}$$

Here $f_0, f_1, \dots, f_4$ and g_1, g_2 denote unknown complex variables and $\alpha_0, \dots, \alpha_4$ are the parameters satisfying the condition:

$$\alpha_0+\alpha_1+2\alpha_2+\alpha_3+\alpha_4=1.$$

Proposition 0.1. *This system has the following invariant divisors*:

Key words and phrases. Affine Weyl group, birational symmetries, coupled Painlevé systems.
2000 Mathematics Subject Classification Numbers. 34M55, 34M45, 58F05, 32S65.

invariant divisors	parameter's relation
$f_0 := 0$	$\alpha_0 = 0$
$f_1 := 0$	$\alpha_1 = 0$
$f_2 := 0$	$\alpha_2 = 0$
$f_3 := 0$	$\alpha_3 = 0$
$f_4 := 0$	$\alpha_4 = 0$

Theorem 0.1. *This system is invariant under the transformations* $s_0, \dots, s_4$ *defined as follows*: *with the notation* $(*) := (f_0, f_1, \dots, f_4, g_1, g_2; \alpha_0, \alpha_1, \dots, \alpha_4)$,

$$\begin{aligned}
&s_0 : (*) \rightarrow \left(f_0, f_1, f_2 + \frac{\alpha_0 g_2}{f_0}, f_3, f_4, g_1 + \frac{\alpha_0}{f_0}, g_2; -\alpha_0, \alpha_1, \alpha_2 + \alpha_0, \alpha_3, \alpha_4\right),\\
&s_1 : (*) \rightarrow \left(f_0, f_1, f_2 + \frac{\alpha_1 g_2}{f_1}, f_3, f_4, g_1 + \frac{\alpha_1}{f_1}, g_2; \alpha_0, -\alpha_1, \alpha_2 + \alpha_1, \alpha_3, \alpha_4\right),\\
&s_2 : (*) \rightarrow \left(f_0 - \frac{\alpha_2 g_2}{f_2}, f_1 - \frac{\alpha_2 g_2}{f_2}, f_2, f_3 - \frac{\alpha_2 g_1}{f_2}, f_4 - \frac{\alpha_2 g_1}{f_2}, g_1, g_2;\right.\\
&\qquad \left.\alpha_0 + \alpha_2, \alpha_1 + \alpha_2, -\alpha_2, \alpha_3 + \alpha_2, \alpha_4 + \alpha_2\right),\\
&s_3 : (*) \rightarrow \left(f_0, f_1, f_2 + \frac{\alpha_3 g_1}{f_3}, f_3, f_4, g_1, g_2 + \frac{\alpha_3}{f_3}; \alpha_0, \alpha_1, \alpha_2 + \alpha_3, -\alpha_3, \alpha_4\right),\\
&s_4 : (*) \rightarrow \left(f_0, f_1, f_2 + \frac{\alpha_4 g_1}{f_4}, f_3, f_4, g_1, g_2 + \frac{\alpha_4}{f_4}; \alpha_0, \alpha_1, \alpha_2 + \alpha_4, \alpha_3, -\alpha_4\right).
\end{aligned} \tag{2}$$

Theorem 0.2. *This system has two first integrals*:

$$\frac{d(f_0 - f_1)}{dt} = \frac{d(f_3 - f_4)}{dt} = 0, \quad \frac{d(f_2 - g_1 g_2)}{dt} = f_2 - g_1 g_2.$$

From this, we have

$$f_0 = f_1 - 1, \quad f_3 = f_4 - 1, \quad f_2 - g_1 g_2 = e^{(t+c)}.$$

Here we set

$$t + c = logT, \quad x := g_1,\ y := f_1,\ z := g_2,\ w := f_4,$$

then we obtain a 4-parameter family of coupled Painlevé III systems in dimension four with affine Weyl group symmetry of type $D_4^{(1)}$ explicitly given by

$$\begin{cases}
\dfrac{dx}{dT} = \dfrac{2x^2y - x^2 + (\alpha_0 + \alpha_1)x}{T} - 1 + 2w,\\
\dfrac{dy}{dT} = \dfrac{-2xy^2 + 2xy - (\alpha_0 + \alpha_1)y + \alpha_1}{T},\\
\dfrac{dz}{dT} = \dfrac{2z^2w - z^2 + (\alpha_3 + \alpha_4)z}{T} - 1 + 2y,\\
\dfrac{dw}{dT} = \dfrac{-2zw^2 + 2zw - (\alpha_3 + \alpha_4)w + \alpha_4}{T}
\end{cases} \tag{3}$$

with the Hamiltonian

$$\begin{aligned}
H = &\frac{x^2y^2 - x^2y + (\alpha_0 + \alpha_1)xy - \alpha_1 x}{T} - y\\
&+ \frac{z^2w^2 - z^2w + (\alpha_3 + \alpha_4)zw - \alpha_4 z}{T} - w + 2yw.
\end{aligned} \tag{4}$$



FIGURE 1. The transformations s_i satisfy the relations: $s_i^2=1$ $(i=0,1,2,3,4)$, $(s_0s_1)^2=(s_0s_3)^2=(s_0s_4)^2=(s_1s_3)^2=(s_1s_4)^2=(s_3s_4)^2=1$, $(s_0s_2)^3=(s_1s_2)^3=(s_3s_2)^3=(s_4s_2)^3=1$.

Theorem 0.3. *This system is invariant under the transformations* $s_0,\dots,s_4,\pi_1,\pi_2,\pi_3$ *defined as follows*: *with the notation* $(*):=(x,y,z,w,T;\alpha_0,\alpha_1,\alpha_2,\alpha_3,\alpha_4)$,

$$\begin{aligned}
&s_0:(*)\rightarrow(x+\frac{\alpha_0}{y-1},y,z,w,T;-\alpha_0,\alpha_1,\alpha_2+\alpha_0,\alpha_3,\alpha_4),\\
&s_1:(*)\rightarrow(x+\frac{\alpha_1}{y},y,z,w,T;\alpha_0,-\alpha_1,\alpha_2+\alpha_1,\alpha_3,\alpha_4),\\
&s_2:(*)\rightarrow(x,y-\frac{\alpha_2 z}{xz+T},z,w-\frac{\alpha_2 x}{xz+T},T;\\
&\qquad\qquad\alpha_0+\alpha_2,\alpha_1+\alpha_2,-\alpha_2,\alpha_3+\alpha_2,\alpha_4+\alpha_2),\\
&s_3:(*)\rightarrow(x,y,z+\frac{\alpha_3}{w-1},w,T;\alpha_0,\alpha_1,\alpha_2+\alpha_3,-\alpha_3,\alpha_4),\\
&s_4:(*)\rightarrow(x,y,z+\frac{\alpha_4}{w},w,T;\alpha_0,\alpha_1,\alpha_2+\alpha_4,\alpha_3,-\alpha_4),\\
&\pi_1:(*)\rightarrow(-x,1-y,z,w,-T;\alpha_1,\alpha_0,\alpha_2,\alpha_3,\alpha_4),\\
&\pi_2:(*)\rightarrow(x,y,-z,1-w,-T;\alpha_0,\alpha_1,\alpha_2,\alpha_4,\alpha_3),\\
&\pi_3:(*)\rightarrow(z,w,x,y,T;\alpha_3,\alpha_4,\alpha_2,\alpha_0,\alpha_1).
\end{aligned}\tag{5}$$

REFERENCES

[1] K. Fuji and T. Suzuki, *The sixth Painlevé equation arising from $D_4^{(1)}$ hierarchy*, J. Phys. A: Math. Gen. **39** (2006), 12073–12082.

[2] T. Matano, A. Matumiya and K. Takano, *On some Hamiltonian structures of Painlevé systems, II*, J. Math. Soc. Japan **51** (1999), 843–866.

[3] M. Noumi and Y. Yamada, *Affine Weyl Groups, Discrete Dynamical Systems and Painlevè Equations*, Comm Math Phys. **199** (1998), 281–295.

[4] M. Noumi, K. Takano and Y. Yamada, *Bäcklund transformations and the manifolds of Painlevè systems*, Funkcial. Ekvac. **45** (2002), 237–258.

[5] M. Noumi, *Painlevé Equations Through Symmetry*, American Mathematical Society, **223** (2004).

[6] Y. Sasano, *Coupled Painlevé V systems in dimension* 4 , Funkcial. Ekvac. **49** (2006), 133–161.

[7] Y. Sasano, *The phase space of coupled Painlevé III system in dimension four*, to appear in Kyoto Journal.

[8] Y. Sasano, *Coupled Painlevé II systems in dimension four and the systems of type $A_4^{(1)}$*, Tohoku. Math. J. **58** (2006), 529–548.

[9] Y. Sasano, *Higher order Painlevé equations of type* ${D_l}^{(1)}$, RIMS Kokyuroku **1473** (2006), 143–163.

[10] Y. Sasano, *Coupled Painlevé VI systems in dimension four with affine Weyl group symmetry of types* $B_6^{(1)}, D_6^{(1)}$ *and* $D_7^{(2)}$, submitted to Nagoya Journal.

[11] Y. Sasano, *Four-dimensional Painlevé systems of types* $D_5^{(1)}$ *and* $B_4^{(1)}$, submitted to Nagoya Journal.

[12] Y. Sasano and Y. Yamada, *Symmetry and holomorphy of Painlevé type systems*, RIMS Kokyuroku **B2** (2007), 215–225.

[13] T. Shioda and K. Takano, *On some Hamiltonian structures of Painlevé systems I*, Funkcial. Ekvac. **40** (1997), 271–291.

[14] M. Suzuki, N. Tahara and K. Takano, *Hierarchy of Bäcklund transformation groups of the Painlevé equations*, J. Math. Soc. Japan **56** (2004), 1221–1232.

[15] N. Tahara, *An augmentation of the phase space of the system of type* $A_4^{(1)}$, Kyushu J. Math. **58** (2004), 393–425.

Graduate School of Mathematical Sciences, The University of Tokyo, Tokyo 153-8914, Japan.

E-mail address: `sasano@ms.u-tokyo.ac.jp`